

14 Konverteringsguide opioider

Morfin		Oxikodon		Hydro- morfon (Palladon) fr.a. till pump		Fentanyl
Dygnsdos mg	Dygnsdos mg	Dygnsdos mg	Dygnsdos mg	Dygnsdos mg	Dygnsdos mg	µg/h
po	sc/iv	po	sc/iv	po	sc/iv	Trans- dermalt
20	7-10	10	7			12
40	15-20	20	15	4-8	2-4	12
60	20-30	30	20	8-12	4-6	25
80	30-40	40	30	12-16	6-8	25
100	35-50	50	35	14-20	7-10	37
120	45-60	60	45	18-24	9-12	50
160	60-80	80	60	24-32	12-16	50
220	80-110	110	80	32-44	16-22	75
320	120-160	160	120	48-64	24-32	100
400	150-200	200	150	60-80	30-40	125
500	185-250	250	185	74-100	37-50	150
580	215-290	290	215	86-116	43-58	175
680	255-340	340	255	102-136	51-68	200
760	285-380	380	285	114-152	57-76	225
860	320-430	430	320	128-172	64-86	250
940	350-470	470	350	140-188	70-94	275
1040	390-520	520	390	156-208	78-104	300

Enl. C. Mucchiano m fl.

Tänk på:

- Varje patient skall dositeras individuellt.
- Försiktig konvertering vid högre doser.
- Vid smärtgenombrott ges en extradosa som utgör ca. 1/6 av dygnsdosen.